

Vaccination Tips

Thank you for choosing the kinetics vaccination services. In order to make everyone feel more at ease in vaccination, please read the following vaccine precautions. If you have any questions, please feel free to contact us.



General Vaccine Precautions

- 1 For all vaccination services, please make an appointment by phone or online at least one week prior to the vaccination session. (if the appointment is on Saturday or close to public holidays, please make the appointment two/three weeks in advance)
- 2 All online reservation must be registered at least three working days prior to the appointment, all reservations without our confirmation will not be accepted.
- 3 If you have physical discomfort or fever on the day of injection, we recommend to postpone injection until you recover.
- 4 The majority of the vaccine can be vaccinated together on the same day (different regions of the body, eg. left/right deltoid), such as influenza and pneumococcal vaccines, for more information, please contact us.
- 5 Some of the vaccines are not suitable for women who are planning pregnancy, pregnant or breastfeeding, if you have any doubts, please inform our medical personnel.
- 6 If you have taken any medication (including herbal) before vaccination, please notify our medical personnel, if possible, you can also bring along the medication when inquiry.
- 7 Some of the vaccines have the scheduled up to 18 months, please remember to complete the vaccination for optimal protection. Our center provides SMS alert service, SMS will be sent 1-2 days prior to the vaccination to scheduled.

Children Vaccine Precautions

- 1 Pay more attention to your child's physical condition a few days prior to vaccination, if there is any discomfort or fever, it is recommended to wait until your child recover to get the vaccine. If you have any questions, please contact us.
- 2 The parent please keep the Immunisation record (injection card) properly and bring along the card for record for every visits.
- 3 After vaccination, children may have fever, gastrointestinal discomfort, swelling around the injection site or other kinds of physical discomfort. With proper post-vaccination care, parents should not worry too much. If there is any abnormal situation, it is recommended to seek medical assistance.
- 4 For children, fevers after vaccination in general are relatively minor, with the body temperature around 38 degrees Celsius, after live attenuated vaccine as the fever may delayed till 14-21 days, parents should pay more attention.
- 5 After the inoculation of oral rotavirus vaccine, some children will experience stomach upset or loose stools, parents should not worry too much. If you have any questions, please contact us.